



Hormonal effects on women's facial masculinity preferences: The influence of pregnancy, post-partum, and hormonal contraceptive use



Kelly D. Cobey*, Anthony C. Little, S. Craig Roberts

School of Natural Sciences, Department of Psychology, University of Stirling, Scotland FK9 4LA, United Kingdom

ARTICLE INFO

Article history:

Received 8 July 2014

Accepted 6 November 2014

Available online 25 November 2014

Keywords:

Face perception

Sexual dimorphism

Pregnancy

Post-partum

Hormonal contraception

ABSTRACT

Here, we investigate changes in women's facial masculinity preferences across pregnancy and the post-partum period. The majority of previous research demonstrating changes in women's masculinity preferences has examined the impact of hormonal variation across the female menstrual cycle. Hormonal changes experienced during pregnancy and the post-partum period, critical periods in women's reproductive life histories, are considerably more extreme than the variation that occurs across the menstrual cycle, suggesting that differences in preferences may also be displayed during these times. We find that women's preference for masculinity in men's faces, but not women's faces, decreases in the post-partum period relative to pregnancy. Furthermore, when compared to a sample of nulliparous control participants, post-partum participants showed different masculinity preferences compared with women who were using hormonal contraception, with the direction of this difference dependent upon the sex of the face assessed.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

A large number of studies indicate that women's preference for masculinity in male faces is moderated by their menstrual cycle stage. For example, in the late follicular phase of the cycle, when oestrogen levels are relatively high and conception risk is highest, women have stronger preferences for facial masculinity than they do in the luteal phase of the cycle when oestrogen and conception risk are low (Little, Jones, & DeBruine, 2008; Little & Jones, 2012; Penton-Voak & Perrett, 2000; Penton-Voak et al., 1999). Increased attraction to masculinity during the late follicular phase has likewise been documented in women's ratings of male voices (Feinberg et al., 2006; Puts, 2005), bodies (Little, Jones, & Burriss, 2007), and masculine characteristics in point-light displays of biological motion (Provost, Troje, & Quinsey, 2008, but see also Peters, Simmons, & Rhodes, 2009). An increased preference for related traits, such as dominant behavioural displays (Gangestad, Simpson, Cousins, Garver-Apgar, & Christensen, 2004), personality traits associated with dominance (Lukaszewski & Roney, 2009), the odour of dominant men (Havlicek, Roberts, & Flegr, 2005), and men's height (Pawlowski & Jasienska, 2005), have also been shown to occur in the follicular phase of the cycle.

One possible explanation for shifts in judgements of masculinity and related traits across the menstrual cycle is that at mid-cycle, when conception risk is highest, cues to a man's condition or genetic quality are especially pertinent for women in order to make an adaptive mate choice. Masculine-faced men are thought to possess 'good genes' and to be healthier than relatively feminine-faced men (Thornhill & Gangestad, 2006), suggesting that a selective preference for masculinity when conception is possible could increase reproductive success. Indeed, masculinity has long been suggested to be an indicator of male quality (Folstad & Karter, 1992; Thornhill & Gangestad, 1999), although there is limited evidence for this proposition (e.g. Scott, Clark, Boothroyd, & Penton-Voak, 2013). Nonetheless, masculine faces are not associated with traits that are desirable in a long term partner, such as investment (Boothroyd, Jones, Burt, & Perrett, 2007), meaning that a general preference for masculinity, even if it does reflect underlying genetic quality, may not be desirable. As such, cycle-dependent changes in line with a woman's conception risk, or current reproductive condition, could carry adaptive benefits.

Irrespective of the logic for why such shifts occur, that cyclical changes in women's preferences for masculine traits occur suggests that within-individual variation in judgements, at least in part, are associated with hormonal variation. Indeed, existing work looking at other phases of hormonal transition across the female lifespan appears consistent with this idea. For example, circum-menopausal women show an increased preference for femininity in male and female faces relative to regularly cycling women

* Corresponding author. Tel.: +44 1786 473171.

E-mail address: kelly.cobey@stir.ac.uk (K.D. Cobey).

(Jones, Vukovic, Little, Roberts, & DeBruine, 2011; Little et al., 2010; Vukovic et al., 2009). This finding may be explained by the reduced value placed on adaptive mate choice among circum-menopausal aged women. Specifically, circum-menopausal women may benefit less from choosing a masculine male partner since conception is no longer a relevant concern, and thus the advertised importance of genetic quality in men may be of diminished value. Moreover, increased preference for femininity in female faces may reflect lower levels of intrasexual competition, with women of this age achieving less from rival derogation relative to those of reproductive age. In other related work, Little et al. (2010) showed that post-pubescent girls had stronger preferences for masculinity in male faces than did younger peri-pubescent girls (see also Saxton, DeBruine, Jones, Little, & Roberts (2009), for similar results in both faces and voices). Together, these findings appear to suggest that attraction to masculinity is particularly high at times in which women are reproductively active. Advertisements of male quality may be most salient to women in the reproductive age group, particularly when fertile.

While it is obvious that menarche, the menstrual cycle and the transition to menopause are related to major shifts in women's hormonal profiles, much research remains in determining which hormone, or group of hormones, mediate these changes in judgements of masculinity. Work by Welling et al. (2007) has previously documented a positive association between women's salivary testosterone and preference for male facial masculinity. This finding is consistent with much of the literature outlined above. For example, while effects are mixed, some research suggests very moderate testosterone increases occur near to ovulation in the follicular phase of the cycle (e.g. Alexander, Sherwin, Bancroft, & Davidson, 1990; Dabbs, 1990; but see also: Schultheiss, Dargel, & Rohde, 2003; Welling et al., 2007) and indeed this is the time when women express enhanced masculinity preferences. Moreover, post-pubescent girls have higher levels of testosterone than peri-pubescent girls (e.g. Angold, Costello, Erkanli, & Worthman, 1999) and, while testosterone does not show a precipitous drop in the menopause, it does show a gradual age decline, meaning older women are likely to have lower levels of this hormone (Longcope, Franz, Morello, Baker, & Johnston, 1986; Zumof, Strain, Miller, & Israel, 1995). Therefore, the finding that testosterone positively relates to masculinity preferences is consistent with the pattern of results documented across these phases of female hormonal transition. Nonetheless, a range of other hormones, including oestrogen and progesterone, fluctuate across these reproductive life events. Thus, it is possible that several hormones, or interactions among hormones, underpin shifts in masculinity preferences that have been documented across the female lifespan.

The current experiment investigated the impact of pregnancy and the post-partum period on women's preferences for feminized versus masculinized versions of both same-sex and opposite-sex faces. If variation in testosterone or other ovarian hormones contributes to masculinity preferences, pregnancy and the post-partum period may reflect further life stages wherein women's preferences for this trait change. It is known that testosterone levels increase throughout pregnancy and then drop swiftly post-partum (e.g. Buckwalter et al., 1999; Leary, Boyne, Flett, Beilby, & James, 1991). Given the results of Welling et al. (2007), discussed above, this may suggest that women would exhibit decreased preference for masculinity post-partum relative to pregnancy. Moreover, Watkins (2012) showed that masculinity preferences relate to measures of reproductive interest or ambition, and, if this is the case, we might further expect pregnancy and the post-partum to be associated with lower masculinity preferences than those exhibited by non-pregnant women of reproductive age. Notably, regularly cycling women have been shown to prefer greater levels of masculinity than women who use hormonal contraception (e.g.

Jones et al., 2005; Little, Burriss, Petrie, Jones, & Roberts, 2013). Pre-existing differences between oral contraceptive users and regularly cycling women may result in divergent differences when these groups' preferences are compared to

3. Results

3.1. Preferences for male faces

We first computed the mean masculinity preference scores for the ten male and female face pairs for participants when pregnant and in the post-partum period. We coded the selection of the feminine face in each pair as ‘zero’ and the selection of the masculine face as ‘one’. Thus, participants were assigned an overall score ranging from zero to ten, with higher scores indicating a greater preference for masculine faces. We similarly calculated the masculinity preferences among our group of nulliparous participants. Paired *t*-tests indicated that women preferred higher rates of masculinity in male faces than in female faces, a difference which was significant both when participants were pregnant ($t = 4.67$, $df = 25$, $p < .001$) and in the post-partum phase ($t = 2.63$, $df = 25$, $p = .015$). Note that the sample for these analyses was somewhat reduced since there was missing data from 2 women’s judgments of the female faces. This difference was also noted among our control sample of nulliparous women who were regularly cycling ($t = 4.27$, $df = 32$, $p < .001$) and among those who were using hormonal contraception ($t = 9.88$, $df = 41$, $p < .001$). Non-parametric tests of these differences revealed similar and significant effects.

To test the impact of transitioning from pregnancy to the post-partum period on masculinity preferences, we conducted a linear-mixed model with reproductive state (pregnancy or post-partum) as the within-participant fixed factor and masculinity preference score obtained from the male faces as the dependent variable. The results of this test indicated a significant main effect of measurement time ($F = 7.63$, $p = .01$). Participants indicated a greater preference for masculine male faces during pregnancy ($M = 6.15$, $S.E. = .49$) relative to the post-partum phase ($M = 4.89$, $S.E. = .49$) (Fig. 1). We then tested the main effect of participant age by adding this variable as a covariate to the model. Age did not have a significant main effect ($F = .001$, $p = .98$) on the model, and the main effect of reproductive state remained significant ($F = 7.61$, $p = .01$). Since previous research indicates differences in masculinity preferences related to use of hormonal contraception (e.g. Jones et al., 2005; Little, Burriss, Petrie, Jones, & Roberts, 2013), we then re-ran this analysis adding whether or not the participant had begun to use hormonal contraception since the birth (yes/no, $N = 9$ had begun hormonal contraception) as a further fixed factor. In this model, the main effect of measurement time remained significant ($F = 7.01$, $p = .01$) and there was a main effect of contraceptive use in the post-partum phase ($F = 8.55$, $p = .01$), with women who used hormonal




Fig. 1. Mean (+S.E.) masculinity preference scores for assessments of male faces. Participants preferred significantly higher levels of masculinity in male faces when pregnant, compared to the post-partum period.

contraception preferring higher levels of masculinity. However, the interaction between measurement time and contraceptive use was non-significant ($F = .44$, $p = .51$). Thus, irrespective of use or non-use of hormonal contraception in the post-partum phase, women preferred lower levels of masculinity in male faces at this time relative to pregnancy.

We then examined how male facial masculinity preferences during pregnancy and post-partum compared to preferences reported by our nulliparous control sample. Since some women in our nulliparous group were using hormonal contraception, we compared regularly cycling and hormonal contraceptive using women as distinct control groups. When pregnant, our participants gave scores that fell between those given by regularly cycling and contraceptive using control participants, but were not significantly different from either (Mean difference = 1.14, $S.E. = .74$, $p = .13$; Mean difference = $-.69$, $S.E. = .70$, $p = .33$). In contrast to previous work, women using hormonal contraceptives preferred higher levels of masculinity in male faces than women who were regularly cycling (mean difference = 1.83, $S.E. = .67$, $p < .01$). Women’s post-partum scores were significantly lower than those given by women who used hormonal contraception (mean difference = -1.94 , $S.E. = .70$, $p < .01$), but did not differ from women who were regularly cycling (mean difference = $-.11$, $S.E. = .74$, $p = .89$) (Fig. 2a).




Fig. 2. Masculinity preference scores for assessments of (a) male faces and (b) female faces. Shading reflects distinct groups of participants; note that the grey bars are repeated measures from the same individuals. In assessing male faces, women using hormonal contraception showed a greater preference for masculinity than regularly cycling women or women in the post-partum. In assessing female faces, women using hormonal contraception and women who were regularly cycling showed a lower preference for masculinity than pregnant and post-partum women.

3.2. Judgments of female faces

Next, we re-ran the same tests described above, this time testing changes in preferences for masculinity in women's faces. The results of the linear-mixed model testing within-person shifts in masculinity preference from pregnancy to the post-partum phase indicated that there was no significant effect of measurement time on masculinity judgement ($F = .009, p = .92$). When participant age was added as covariate to the model it remained non-significant ($F = .01, p = .92$) and age did not have a main effect ($F = .46, p = .50$). Moreover, when we split the model based on women's hormonal contraceptive use status in the post-partum phase, the models were non-significant (non-users: $F = 1.05, p = .35$, Hormonal contraceptive users: $F = .87, p = .36$). Taken together, these results suggest that between-individual judgements of masculinity in female faces do not vary significantly across the transition from pregnancy to the post-partum (Fig. 1).

When we examined how pregnant and post-partum attractiveness judgements for masculinity in female faces differed from our control group of nulliparous women's scores, we found that pregnant women judged higher levels of masculinity as more attractive than women who were using hormonal contraception (mean difference = 1.60, S.E. = .59, $p < .01$). The difference between our pregnant participants judgements and that of the regularly cycling control group was non-significant (mean difference = .89, S.E. = .63, $p = .16$). Similarly, post-partum scores were significantly higher than scores given by hormonal contraceptive users (mean difference = 1.52, S.E. = .60, $p = .01$), but did not differ from women who were regularly cycling (mean difference = .81, S.E. = .63, $p = .20$). There was no difference in preferences between our control group of nulliparous women using hormonal contraception and nulliparous regularly cycling women (mean difference = $-.71$, S.E. = .55, $p = .20$) (See Fig. 2b).

4. Discussion

The current study demonstrated a prospective shift in preference for masculinity in male faces from pregnancy to the post-partum phase measured within-subjects. Specifically, participants preferred relatively less masculine male faces in the weeks following birth compared to while they were pregnant. This finding is consistent with the idea that masculinity preferences for male faces are higher when testosterone levels are higher (Welling et al., 2007). In contrast, preferences for masculinity in female faces were unchanged between the two measurement times. That shifts in preferences for masculinity are specific to judgments of male faces appears to suggest that effects of hormonal variation between the two measurement times impacts partner preferences specifically, rather than reflecting changes in general face preferences. When one considers the high demands associated with rearing newborns (e.g. Clutton-Brock, 1991), reduced attraction to mating relevant cues in the post-partum phase may be considered as an adaptive means to promote maternal investment. Moreover, femininity in male faces reflects cues to pro-social behaviour (Watkins, DeBruine, Little, & Jones, 2012), suggesting it may additionally be adaptive for women with young infants, a time when investment is important, to prefer men displaying lower levels of masculinity since this may suggest they are likely to be more investing.

Our participants' masculinity judgements when pregnant and in the post-partum phase were also compared with a control sample of nulliparous women's judgments. We found that when judging male faces, women in the post-partum period preferred lower levels of masculinity than our hormonal contraceptive group. However, the reverse occurred when these same women judged the attractiveness of female faces: our post-partum group preferred

significantly higher levels of masculinity in female faces than our hormonal contraceptive using group. This result appears to suggest that endocrine changes associated with the post-partum phase influence facial masculinity preferences distinctly for the assessments of male and female stimuli. Again, this finding may suggest that indicators of male quality, or cues to rival females in the environment, are less salient or desired given the demands associated with motherhood.

Finally, our results indicated that women who were using hormonal contraception preferred greater levels of masculinity in male faces compared to women who were regularly cycling, but this difference was not evident for judgments of female faces. This finding is in contrast to previous work on this topic; however, it should be noted that the between-subjects comparison used herein is not as powerful as the existing within-person tests showing the opposite outcome (Little et al., 2013). One factor which may also contribute to our conflicting result is that we did not account for the cycle phase among our regularly cycling women. Moreover, our participants that were taking hormonal contraception were not necessarily using the same formulation, giving rise to the possibility that different formulations have different effects, or effects of differing magnitude (e.g. Cobey, Pollet, Roberts, & Buunk, 2011; Welling, Puts, Roberts, Little, & Burriss, 2012), which could complicate the comparison made. Naturally, it is also possible that differences in masculinity preferences observed between users and non-users are the product of baseline differences (e.g. in reproductive ambition or relationship status) between the groups which were

lifespan, namely pregnancy and the post-partum phase, that influence women's facial judgements. This work benefited from the use of a within-subjects design to test the impact of transitioning from pregnancy to the post-partum phase, and the consideration of these preferences relative to two control groups. Furthermore, the use of both male and female facial stimuli allows for conclusions to be drawn with respect to the specificity versus generality of the finding (i.e. is it a mating-relevant shift or a general shift in face preference?). As noted, one limitation of the current work is that we did not account for the cycle stage of our nulliparous control group of regularly cycling women. However, given that we detected significant differences between groups in our comparisons of pregnant women, post-partum women, and the control groups, more subtle within-individual variation may be less relevant. It is also worth acknowledging that the control groups in our study, namely those women who were regularly cycling or using hormonal contraception, completed the facial rating task on just one occasion. In contrast, our experimental group completed the task twice: once during pregnancy and then again during the post-partum phase. While we feel it is unlikely, it is possible that there is a design confound in that we did not conduct a second measure among our control participants, and we were unable to randomize the order of the two naturally occurring phases in our experimental group. Furthermore, since previous work has shown that a woman's combined oral contraceptive pill dosage can also impact mating psychology (Cobey et al., 2011; Welling et al., 2012), future work could also consider preferences of women on different types and formulations of hormonal contraception. Examining the hormone, or group of hormones, responsible for the shifts documented here will certainly be a fruitful area for further research. Testosterone appears to be an obvious hormonal candidate (Welling et al., 2007); however, other hormones that are known to change post-partum (e.g., oestrogen, progesterone, prolactin, oxytocin), or interactions of these hormones, should also be incorporated in future studies.

Acknowledgments

We thank all our participants. SCR was supported by a British Academy Mid-Career Fellowship.

References

- Alexander, G. M., Sherwin, B. B., Bancroft, J., & Davidson, D. W. (1990). Testosterone and sexual behavior in oral contraceptive users and nonusers: A prospective study. *Hormones and Behavior*, 24(3), 388–402.
- Angold, A., Costello, E. J., Erkanli, A., & Worthman, C. M. (1999). Pubertal changes in hormone levels and depression in girls. *Psychological Medicine*, 29(5), 1043–1053. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10576297>
- Benson, P. J., & Perrett, D. I. (1992). *Extracting prototypical facial images from exemplars*, 22, 257–262.
- Berg, S. J., & Wynne-Edwards, K. E. (2001). Changes in testosterone, cortisol, and estradiol levels in men becoming fathers. *Mayo Clinic Proceedings*, 76(6), 582–592. <http://dx.doi.org/10.4065/76.6.582>
- Boothroyd, L. G., Jones, B. C., Burt, D. M., & Perrett, D. I. (2007). Partner characteristics associated with masculinity, health and maturity in male faces. *Personality and Individual Differences*, 43(5), 1161–1173. <http://dx.doi.org/10.1016/j.paid.2007.03.008>
- Buckwalter, J. G., Stanczyk, F. Z., McCleary, C. A., Bluestein, B. W., Buckwalter, D. K., Rankin, K. P., et al. (1999). Pregnancy, the postpartum, and steroid hormones: Effects on cognition and mood. *Psychoneuroendocrinology*, 24(1), 69–84. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10098220>
- Clutton-Brock, T. H. (1991). *The evolution of parental care*. Princeton, NJ: Princeton University Press.
- Cobey, K. D., Pollet, T. V., Roberts, S. C., & Buunk, A. P. (2011). Hormonal birth control use and relationship jealousy: Evidence for estrogen dosage effects. *Personality and Individual Differences*, 50(2), 315–317. <http://dx.doi.org/10.1016/j.paid.2010.09.012>
- Dabbs, J. M. (1990). Salivary testosterone measurements: Reliability across hours, days, and weeks. *Physiology & Behavior*, 48(1), 83–86. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/2236282>
- DeBruine, L., Jones, B. C., Frederick, D. A., Haselton, M. G., Penton-voak, I. S., & Perrett, D. (2010). Evidence for menstrual cycle shifts in women's preferences for masculinity: A response to Harris (in press) "Menstrual cycle and facial preferences reconsidered". *Evolutionary Psychology*, 8(4), 768–775.
- Feinberg, D. R., Jones, B. C., Law Smith, M. J., Moore, F. R., DeBruine, L. M., Cornwell, R. E., et al. (2006). Menstrual cycle, trait estrogen level, and masculinity preferences in the human voice. *Hormones and Behavior*, 49(2), 215–222. <http://dx.doi.org/10.1016/j.yhbeh.2005.07.004>
- Feinberg, D. R., DeBruine, L. M., Jones, B. C., & Little, A. C. (2008). Correlated preferences for men's facial and vocal masculinity. *Evolution and Human Behavior*, 29(4), 233–241. <http://dx.doi.org/10.1016/j.evolhumbehav.2007.12.008>
- Folstad, I., & Karter, A. J. (1992). Parasites, bright males, and the immunocompetence handicap. *American Naturalist*, 139(3), 603–622.
- Gangestad, S. W., Simpson, J. A., Cousins, A. J., Garver-Appar, C. E., & Christensen, P. N. (2004). Women's preferences for male behavioral displays change across the menstrual cycle. *Psychological Science*, 15(3), 203–207. <http://dx.doi.org/10.1111/j.0956-7976.2004.01503010.x>
- Gettler, L. T., McDade, T. W., Feranil, A. B., & Kuzawa, C. W. (2011). Longitudinal evidence that fatherhood decreases testosterone in human males. *Proceedings of the National Academy of Sciences*, 108(39), 16194–16199. <http://dx.doi.org/10.1073/pnas.1105403108>
- Havlicek, J., Roberts, S. C., & Flegr, J. (2005). Women's preference for dominant male odour: Effects of menstrual cycle and relationship status. *Biology Letters*, 1(3), 256–259. <http://dx.doi.org/10.1098/rsbl.2005.0332>
- Jones, B. C., Perrett, D. I., Little, A. C., Boothroyd, L., Cornwell, R. E., Feinberg, D. R., et al. (2005). Menstrual cycle, pregnancy and oral contraceptive use alter attraction to apparent health in faces. *Proceedings of the Royal Society B-Biological Sciences*, 272(1561), 347–354. <http://dx.doi.org/10.1098/rspb.2004.2962>
- Jones, B. C., Vukovic, J., Little, A. C., Roberts, S. C., & DeBruine, L. M. (2011). Circum-menopausal changes in women's preferences for sexually dimorphic shape cues in peer-aged faces. *Biological Psychology*, 87(3), 453–455. <http://dx.doi.org/10.1016/j.biopsycho.2011.04.004>
- Kuzawa, C. W., Gettler, L. T., Huang, Y., & McDade, T. W. (2010). Mothers have lower testosterone than non-mothers: Evidence from the Philippines. *Hormones and Behavior*, 57(4–5), 441–447. <http://dx.doi.org/10.1016/j.yhbeh.2010.01.014>
- Leary, P. O., Boyne, P., Flett, P., Beilby, J., & James, I. (1991). Longitudinal assessment of pregnancy in reproductive hormones during normal pregnancy. *Clinical Chemistry*, 37(5), 667–672.
- Little, A. C., Burriss, R. P., Petrie, M., Jones, B. C., & Roberts, S. C. (2013). Oral contraceptive use in women changes preferences for male facial masculinity and is associated with partner facial masculinity. *Psychoneuroendocrinology*, 38(9), 1777–1785. <http://dx.doi.org/10.1016/j.psyneuen.2013.02.014>
- Little, A. C., Cohen, D. L., Jones, B. C., & Belsky, J. (2006). Human preferences for facial masculinity change with relationship type and environmental harshness. *Behavioral Ecology and Sociobiology*, 61(6), 967–973. <http://dx.doi.org/10.1007/s00265-006-0325-7>
- Little, A. C., & Jones, B. C. (2012). Variation in facial masculinity and symmetry preferences across the menstrual cycle is moderated by relationship context. *Psychoneuroendocrinology*, 37(7), 999–1008. <http://dx.doi.org/10.1016/j.psyneuen.2011.11.007>
- Little, A. C., Jones, B. C., & Burriss, R. P. (2007). Preferences for masculinity in male bodies change across the menstrual cycle. *Hormones and Behavior*, 51(5), 633–639. <http://dx.doi.org/10.1016/j.yhbeh.2007.03.006>
- Little, A. C., Jones, B. C., & DeBruine, L. M. (2008). Preferences for variation in masculinity in real male faces change across the menstrual cycle: Women prefer more masculine faces when they are more fertile. *Personality and Individual Differences*, 45(6), 478–482. <http://dx.doi.org/10.1016/j.paid.2008.05.024>
- Little, A. C., Saxton, T. K., Roberts, S. C., Jones, B. C., DeBruine, L. M., Vukovic, J., et al. (2010). Women's preferences for masculinity in male faces are highest during reproductive age range and lower around puberty and post-menopause. *Psychoneuroendocrinology*, 35(6), 912–920. <http://dx.doi.org/10.1016/j.psyneuen.2009.12.006>
- Longcope, C., Franz, C., Morello, C., Baker, R., & Johnston, C. C. (1986). Steroid and gonadotropin levels in women during the peri-menopausal years. *Maturitas*, 8(3), 189–196.
- Lukaszewski, A. W., & Roney, J. R. (2009). Estimated hormones predict women's mate preferences for dominant personality traits. *Personality and Individual Differences*, 47(3), 191–196. <http://dx.doi.org/10.1016/j.paid.2009.02.019>
- Pawlowski, B., & Jasienska, G. (2005). Women's preferences for sexual dimorphism in height depend on menstrual cycle phase and expected duration of relationship. *Biological Psychology*, 70(1), 38–43. <http://dx.doi.org/10.1016/j.biopsycho.2005.02.002>
- Penton-Voak, I. S., & Perrett, D. I. (2000). Female preference for male faces changes cyclically: Further evidence. *Evolution and Human Behavior*, 21, 39–48.
- Penton-Voak, I. S., Perrett, D. I., Castles, D. L., Kobayashi, T., Burt, D. M., Murray, L. K., et al. (1999). Menstrual cycle alters face preference. *Nature*, 399(6738), 741–742. <http://dx.doi.org/10.1038/21557>
- Peters, M., Simmons, L. W., & Rhodes, G. (2009). Preferences across the menstrual cycle for masculinity and symmetry in photographs of male faces and bodies. *PLoS ONE*, 4(1), e4138. <http://dx.doi.org/10.1371/journal.pone.0004138>
- Pollet, T. V., Cobey, K. D., & van der Meij, L. (2013). Testosterone levels are negatively associated with childlessness in males, but positively related to offspring count in fathers. *PLoS ONE*, 8(4), e60018. <http://dx.doi.org/10.1371/journal.pone.0060018>
- Provost, M., Troje, N., & Quinsey, V. (2008). Short-term mating strategies and attraction to masculinity in point-light walkers. *Evolution and Human Behavior*, 29(1), 65–69. <http://dx.doi.org/10.1016/j.evolhumbehav.2007.07.007>

- Puts, D. A. (2005). Mating context and menstrual phase affect women's preferences for male voice pitch. *Evolution and Human Behavior*, 26(5), 388–397. <http://dx.doi.org/10.1016/j.evolhumbehav.2005.03.001>
- Roberts, S. C., Cobey, K. D., Klapilová, K., & Havlíček, J. (2013). An evolutionary approach offers a fresh perspective on the relationship between oral contraception and sexual desire. *Archives of Sexual Behavior*, 42(8), 1369–1375. <http://dx.doi.org/10.1007/s10508-013-0126-9>
- Roberts, S. C., Little, A. C., Burriss, R. P., Cobey, K. D., Klapilova, K., Havlíček, J., et al. (2014). Partner choice, relationship satisfaction, and oral contraception: The congruency hypothesis. *Psychological Science*, 25(7), 1497–1503. <http://dx.doi.org/10.1177/0956797614532295>
- Saxton, T. K., DeBruine, L. M., Jones, B. C., Little, A. C., & Roberts, S. C. (2009). Face and voice attractiveness judgments change during adolescence. *Evolution and Human Behavior*, 30(6), 398–408. <http://dx.doi.org/10.1016/j.evolhumbehav.2009.06.004>
- Schultheiss, O. C., Dargel, A., & Rohde, W. (2003). Implicit motives and gonadal steroid hormones: Effects of menstrual cycle phase, oral contraceptive use, and relationship status. *Hormones and Behavior*, 43(2), 293–301. [http://dx.doi.org/10.1016/S0018-506X\(03\)00003-5](http://dx.doi.org/10.1016/S0018-506X(03)00003-5)
- Scott, I. M. L., Clark, A. P., Boothroyd, L. G., & Penton-Voak, I. S. (2013). Do men's faces really signal heritable immunocompetence? *Behavioral Ecology*, 24(3), 579–589. <http://dx.doi.org/10.1093/beheco/ars092>
- Thornhill, R., & Gangestad, S. W. (1999). Facial attractiveness. *Trends in Cognitive Sciences*, 3(12), 452–460. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10562724>
- Thornhill, R., & Gangestad, S. W. (2006). Facial sexual dimorphism, developmental stability, and susceptibility to disease in men and women. *Evolution and Human Behavior*, 27(2), 131–144. <http://dx.doi.org/10.1016/j.evolhumbehav.2005.06.001>
- Vukovic, J., Jones, B. C., DeBruine, L. M., Little, A. C., Feinberg, D. R., & Welling, L. L. M. (2009). Circum-menopausal effects on women's judgements of facial attractiveness. *Biology Letters*, 5(1), 62–64. <http://dx.doi.org/10.1098/rsbl.2008.0478>
- Watkins, C. D. (2012). Reproductive ambition predicts partnered, but not unpartnered, women's preferences for masculine men. *British Journal of Psychology*, 103(3), 317–329. <http://dx.doi.org/10.1111/j.2044-8295.2011.02076.x>
- Watkins, C. D., DeBruine, L. M., Little, A. C., & Jones, B. C. (2012). Social support influences preferences for feminine facial cues in potential social partners. *Experimental Psychology*, 59(6), 340–347. <http://dx.doi.org/10.1027/1618-3169/a000162>
- Welling, L. L. M., Jones, B. C., DeBruine, L. M., Conway, C. A., Law Smith, M. J., Little, A. C., et al. (2007). Raised salivary testosterone in women is associated with increased attraction to masculine faces. *Hormones and Behavior*, 52(2), 156–161. <http://dx.doi.org/10.1016/j.yhbeh.2007.01.010>
- Welling, L. L. M., Puts, D. A., Roberts, S. C., Little, A. C., & Burriss, R. P. (2012). Hormonal contraceptive use and mate retention behavior in women and their male partners. *Hormones and Behavior*, 61(1), 114–120. <http://dx.doi.org/10.1016/j.yhbeh.2011.10.011>
- Zumof, B., Strain, W., Miller, K., & Israel, B. (1995). Twenty-four-hour mean plasma testosterone concentration declines with age in normal premenopausal women. *Journal of Clinical Endocrinology and Metabolism*, 6, 20–21.